

Many-body effect in spin dephasing in n-typed GaAs quantum wells

M. Q. Weng^{1,2} and M. W. Wu^{1,2,*}

¹Structure Research Laboratory, University of Science & Technology of China, Academia Sinica, Hefei, Anhui, 230026, China

²Department of Physics, University of Science & Technology of China, Hefei, Anhui, 230026, China[†]

(Dated: February 2, 2008)

By constructing and numerically solving the kinetic Bloch equations we perform a many-body study of the spin dephasing due to the D'yakonov-Perel' effect in *n*-typed GaAs (100) quantum wells for high temperatures. In our study, we include the spin-conserving scattering such as the electron-phonon, the electron-nonmagnetic impurity as well as the electron-electron Coulomb scattering into consideration. The dephasing obtained from our theory contains not only that due to the effective spin-flip scattering first proposed by D'yakonov and Perel' [Zh. Eksp. Teor. Fiz. **60**, 1954(1971)[Sov. Phys.-JETP **38**, 1053(1971)]], but also the recently proposed many-body dephasing due to the inhomogeneous broadening provided by the DP term [Wu, J. Supercond.:Incorp. Novel Mechanism **14**, 245 (2001); Wu and Ning, Eur. Phys. J. B **18**, 373 (2000)]. Our result agrees with the experiment data [Malinowski *et al.*, Phys. Rev. B **62**, 13034 (2000)] very well. We further show that in the case we study, the spin dephasing is dominated by the many-body effect.

PACS numbers: 71.10.-w, 67.57.Lm, 72.25.Rb, 73.61.Ey

Spintronics,^{1,2,3} where the spin degree of freedom of electrons is proposed to be used in place of/in addition to the charge degree of freedom to add new features and functionalities in semiconductor devices, has attracted intensive attention recently. One of the important prerequisites for the realization of the device application is to understand the spin dephasing. Experiments have shown long spin lifetimes (up to 100 ns) in *n*-typed semiconductors.^{4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19} In theory, three single-particle spin dephasing mechanisms, *ie.* the Elliot-Yafet (EY) mechanism,^{20,21} the D'yakonov-Perel' (DP) mechanism,²² and the Bir-Aronov-Pikus (BAP) mechanism²³, have been proposed and discussed decades ago.²⁴ All these spin dephasing mechanisms require either spin-flip (SF) or effective SF scattering. Recently, Wu proposed a many-body spin dephasing mechanism where the spin conservation (SC) scattering can also cause spin dephasing. The new mechanism, which is caused by irreversibly disrupting the phases between spin dipoles due to the inhomogeneous broadening together with the SC scattering,^{25,26,27,28,29} has long been overlooked. In semiconductors, the inhomogeneous broadening can be introduced by the energy dependence of the *g*-factor^{28,29,30} and/or the momentum **k**-dependence of the DP term.^{26,27,29} In spin diffusion/transport, it may also come from the different spin precession rates for different moment **k** along the spacial gradient which we discussed closely in our recent papers.^{31,32}

It is understood that for *n*-typed GaAs semiconductors, the DP mechanism is the main spin dephasing mechanism at high temperature. Very recently, we performed a systematic investigation^{33,34,35} of the spin dephasing due to the DP effect in *n*-typed narrow quantum wells for high temperatures (≥ 120 K) under magnetic fields in the Voigt configuration by constructing and numerically solving the many-body kinetic Bloch equations.^{25,26,27,28,29,36} With all of the SC scattering (electron-electron, electron-phonon and electron nonmagnetic impurity scattering)

included in our model, we are thus able to study the spin dephasing under various conditions. The spin dephasing time calculated from our theory contains not only the single particle dephasing, but also the many-body dephasing. Moreover, from this many-body theory, we are able to calculate the spin dephasing with arbitrary initial spin polarization (up to 100 %) which cannot be achieved from earlier single particle theory. Our results show many dramatic many-body effects for both high and low spin polarizations.^{33,34,35} Many many-body effects we predicted, especially those under high spin polarizations, have not yet been measured experimentally. In this communication, we will show evidence of the many-body effect by comparing with the existing experiments with small initial spin polarization.

We start our investigation from an *n*-typed (100) GaAs quantum well (QW) with well width *a*. The growth direction is assumed to be *z*-axis. A moderate magnetic field **B** is applied along the *x* axis. Due to the confinement of the QW, the momentum states along *z* axis are quantized. Therefore the electron states are characterized by a subband index *n* and a two dimensional wave vector **k** = (*k_x*, *k_y*) together with a spin index σ . For the QW with narrow width we study in the present paper, only the lowest subband is populated and the transition to the upper subbands is unimportant. With the DP term included, the Hamiltonian of the electrons in the QW takes the form:

$$H = \sum_{\mathbf{k}\sigma\sigma'} \left\{ \varepsilon_{\mathbf{k}} + [g\mu_B \mathbf{B} + \mathbf{h}(\mathbf{k})] \cdot \frac{\vec{\sigma}_{\sigma\sigma'}}{2} \right\} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma'} + H_I. \quad (1)$$

Here $\varepsilon_{\mathbf{k}} = \mathbf{k}^2/2m^*$ is the energy of electron with wavevector **k** and effective mass *m*^{*}. $\vec{\sigma}$ are the Pauli matrices. The DP term **h**(**k**) can be written as^{37,38}

$$h_x^{\text{BIA}}(\mathbf{k}) = \gamma k_x (k_y^2 - \langle k_z^2 \rangle),$$

$$h_y^{\text{BIA}}(\mathbf{k}) = \gamma k_y (\langle k_z^2 \rangle - k_x^2),$$

$$h_z^{\text{BIA}}(\mathbf{k}) = 0. \quad (2)$$

Here $\langle k_z^2 \rangle$ represents the average of the operator $-(\frac{\partial}{\partial z})^2$ over the electronic state of the lowest subband and is therefore $(\pi/a)^2$. $\gamma = (4/3)(m^*/m_{cv})(1/\sqrt{2m^*E_g})(\eta/\sqrt{1-\eta/3})$ and $\eta = \Delta/(E_g + \Delta)$, in which E_g denotes the band gap; Δ represents the spin-orbit splitting of the valence band; and m_{cv} is a constant close in magnitude to the free electron mass m_0 .³⁹ The interaction Hamiltonian H_I is composed of Coulomb interaction H_{ee} , electron-phonon interaction H_{ph} , as well as electron-impurity scattering H_i . Their expressions can be found in textbooks.^{40,41}

We construct the kinetic Bloch equations by the nonequilibrium Green function method⁴⁰ as follows:

$$\dot{\rho}_{\mathbf{k},\sigma\sigma'} = \dot{\rho}_{\mathbf{k},\sigma\sigma'}|_{\text{coh}} + \dot{\rho}_{\mathbf{k},\sigma\sigma'}|_{\text{scatt}}. \quad (3)$$

Here $\rho_{\mathbf{k}}$ represents the single particle density matrix. The diagonal elements describe the electron distribution functions $\rho_{\mathbf{k},\sigma\sigma} = f_{\mathbf{k}\sigma}$. The off-diagonal elements $\rho_{\mathbf{k},\frac{1}{2}-\frac{1}{2}} \equiv \rho_{\mathbf{k}}$ describe the inter-spin-band polarizations (coherence) of the spin coherence.³⁶ Note that $\rho_{\mathbf{k},-\frac{1}{2}\frac{1}{2}} \equiv \rho_{\mathbf{k},\frac{1}{2}-\frac{1}{2}}^* = \rho_{\mathbf{k}}^*$. Therefore, $f_{\mathbf{k}\pm\frac{1}{2}}$ and $\rho_{\mathbf{k}}$ are the quantities to be determined from Bloch equations.

The coherent parts of the equation of motion for the electron distribution function and the spin coherence are given by

$$\frac{\partial f_{\mathbf{k},\sigma}}{\partial t}|_{\text{coh}} = -2\sigma \{ [g\mu_B B + h_x(\mathbf{k})] \text{Im}\rho_{\mathbf{k}} + h_y(\mathbf{k}) \text{Re}\rho_{\mathbf{k}} \} + 4\sigma \text{Im} \sum_{\mathbf{q}} V_{\mathbf{q}} \rho_{\mathbf{k}+\mathbf{q}}^* \rho_{\mathbf{k}}, \quad (4)$$

$$\frac{\partial \rho_{\mathbf{k}}}{\partial t}|_{\text{coh}} = \frac{1}{2} [ig\mu_B B + ih_x(\mathbf{k}) + h_y(\mathbf{k})] (f_{\mathbf{k}\frac{1}{2}} - f_{\mathbf{k}-\frac{1}{2}}) + i \sum_{\mathbf{q}} V_{\mathbf{q}} [(f_{\mathbf{k}+\mathbf{q}\frac{1}{2}} - f_{\mathbf{k}+\mathbf{q}-\frac{1}{2}}) \rho_{\mathbf{k}} - \rho_{\mathbf{k}+\mathbf{q}} (f_{\mathbf{k}\frac{1}{2}} - f_{\mathbf{k}-\frac{1}{2}})], \quad (5)$$

where $V_{\mathbf{q}} = 4\pi e^2/[\kappa_0(q + q_0)]$ is the 2D Coulomb matrix element under static screening. $q_0 = (e^2 m^*/\kappa_0) \sum_{\sigma} f_{\mathbf{k}=0,\sigma}$ and κ_0 is the static dielectric constant. The first term on the right hand side (RHS) of Eqs. (4) and (5) describes spin precession of electrons under the magnetic field $\mathbf{B}$ as well as the effective magnetic field $\mathbf{h}(\mathbf{k})$ due to the DP effect. The rest terms are the contribution to the spin precession from the HF term of the electron-electron interaction. $\dot{\rho}_{\mathbf{k},\sigma\sigma'}|_{\text{scatt}}$ stands for the scattering term. In our theory, we include all of the SC scattering, *i.e.*, the electron-non-magnetic impurity, electron-phonon as well as the electron-electron scattering. Their expressions are listed in detail in Refs. 33,34,35, and will not be repeated here.

The initial conditions are taken at $t = 0$ as:

$$\rho_{\mathbf{k}}|_{t=0} = 0 \quad (6)$$

$$f_{\mathbf{k}\sigma}|_{t=0} = 1/\{\exp[(\varepsilon_{\mathbf{k}} - \mu_{\sigma})/k_B T] + 1\} \quad (7)$$

where μ_{σ} is the chemical potential for spin σ . The condition $\mu_{\frac{1}{2}} \neq \mu_{-\frac{1}{2}}$ gives rise to the imbalance of the electron densities of the two spin bands. Eqs. (3) through (5) together with the initial conditions Eqs. (6) and (7) comprise the complete set of kinetic Bloch equations of our investigation.

The kinetic Bloch equations form a set of nonlinear equations. The DP term in coherent part of the Bloch equation tends to drive the electron distribution out of equilibrium Fermi distribution, while the scattering term attempts to randomize electrons in $\mathbf{k}$ -space. There-

fore, these two parts are coupled in the Bloch equations and the equations have to be solved self-consistently. This self-consistent solution can be performed numerically through the method that has been laid out in detail in the previous paper on the DP mechanism in 3D systems.²⁵ By numerically solving these coupled equations, we are thus able to study the spin precession between spin-up and -down bands. As discussed in the previous papers,^{36,42} one can obtain the spin dephasing time as the inverse of the slope of the envelope of the incoherently summed spin coherence $\rho(t) = \sum_{\mathbf{k}} |\rho_{\mathbf{k}}|$. This spin dephasing time consists both the single-particle effective-SF-scattering-induced spin dephasing time and the many-body spin dephasing time. We will compare our spin dephasing time with the experiments as well as that predicted by the widely used single particle theory:^{27,43,44}

$$\frac{1}{\tau} = \frac{\int_0^{\infty} dE_k (f_{k\frac{1}{2}} - f_{k-\frac{1}{2}}) \Gamma(k)}{\int_0^{\infty} dE_k (f_{k\frac{1}{2}} - f_{k-\frac{1}{2}})}, \quad (8)$$

with

$$\Gamma(k) = 2\tau_1(k) \left[(\gamma \langle k_z^2 \rangle)^2 k^2 - \frac{1}{2} \gamma \langle k_z^2 \rangle k^4 + \frac{1 + \tau_3(k)/\tau_1(k)}{16} \gamma^2 k^6 \right] \quad (9)$$

$$\tau_n^{-1}(k) = \int_0^{2\pi} \sigma(E_k, \theta) [1 - \cos(n\theta)] d\theta. \quad (10)$$

Here $\sigma(E_k, \theta)$ stands for the scattering cross-section.

We compare the spin dephasing times calculated from our many-body theory in two GaAs QW's for the well width a being 6 nm and 10 nm with those from the experiment by Malinowski *et al.*⁴⁵ In the experiment the spin dephasing times for 15 and 20 nm wells have also been measured. However, for these wider width QW's, one needs more subbands than just the lowest one as we use in the present paper. The electron density is $N_e = 2 \times 10^{11} \text{ cm}^{-2}$ which is the background excitation measured in the experiment for both samples. The material parameters of GaAs for our calculation are tabulated in Table I.⁴⁶ The impurity densities N_i are taken to be $0.1N_e$ for $a = 6 \text{ nm}$ QW and $0.05N_e$ for $a = 10 \text{ nm}$ QW. It is noted that here we use the elastic impurity scattering to simulate also the electron scattering by the interface roughness. Therefore it is understood that the impurity density for the 6 nm QW is larger than that for the 10 nm well as narrower QW suffers more scattering from the interface roughness.

κ_∞	10.8	κ_0	12.9
ω_0	35.4 meV	m^*	$0.067 m_0$
Δ	0.341 eV	E_g	1.55 eV
g	0.44	m_{cv}	$1.25 m_0$

TABLE I: Parameters used in the numerical calculations

In Fig. 1 the spin dephasing time is plotted as a function of the temperature for 10 nm QW (a) and 6 nm QW (b). The corresponding experimental data are plotted as dots. For comparison, the spin dephasing time calculated from the single particle theory Eq. (8) from the *same* parameters is also given as dashed curve in the same figure. It is seen exactly from the figure that our results fit very well with the experiment data. In contrast, the spin dephasing time predicted by the single particle model is about one order of magnitude *slower* than the experiment data. Moreover, it is noticed that when the temperature increases, both our theory and experiment data indicate that the spin dephasing time slightly *increases* with the temperature whereas the single particle model predicates a dramatic *decrease* of the spin dephasing time with the increase of the temperature.

These results clearly demonstrate the important role the many-body effect plays in the spin dephasing. The single particle theory does not count for the many-body dephasing due to the inhomogeneous broadening inherited in the DP term.^{25,26,27,28,29} Therefore, it is easy to understand why the single particle model gives much slower spin dephasing time.

Taking the inhomogeneous broadening effect into account, one can also understand the temperature dependence of the spin dephasing time: as the temperature increases, the inhomogeneous broadening is reduced as the electrons are distributed to the wider $\mathbf{k}$ -states. As the result, the number of electron occupation on each $\mathbf{k}$ state around the Fermi surface is reduced. Consequently




FIG. 1: The spin dephasing time as a function of the temperature for two GaAs quantum wells with different well width: (a) $a=10 \text{ nm}$; (b) $a=6 \text{ nm}$.

the inhomogeneous broadening and therefore the many-body spin dephasing are suppressed. It is further noted that this reduction is mild as a function of the temperature. Therefore, the temperature dependence is quite mild.

In summary, we have shown evidence of the many-body effect in the spin dephasing for n -typed narrow QW's where the spin dephasing is dominated by the many-body effect. It is noted that the small well width not only limits electrons to the lowest subband, but also enhances the DP term as $\langle k_z^2 \rangle = (\pi/a)^2$ in Eq. (2) is greatly enhanced for small width and it is much larger than k_F^2 for the density around 10^{11} cm^{-2} . Consequently the inhomoge-

neous broadening is also greatly enhanced. For wide QW or bulk sample, the DP term is not enhanced. We notice from the experiment that the temperature dependence of the spin dephasing time is similar to what predicted by the single particle theory, *ie.*, the spin dephasing time decreases with the temperature. A full many-body quantum kinetic calculation for the quantitative accurate spin dephasing time is very time consuming and is still missing. Our earlier qualitative calculation for bulk GaAs has

shown the decrease of the spin dephasing time with the temperature.²⁵ A thorough investigation is still underway to find out how large the many-body effect is for those samples.

MWW is supported by the “100 Person Project” of Chinese Academy of Sciences and Natural Science Foundation of China.

* Author to whom correspondence should be addressed; Electronic address: mwwu@ustc.edu.cn

† Mailing Address.

- ¹ M. Ziese and M. J. Thornton, *Spin Electronics* (Springer, Berlin, 2001).
- ² S. A. Wolf, D. D. Awschalom, R. A. Buhrmann, J. M. Daughton, S. von Malnär, M. L. Roukes, A. Y. Chtchelkanova, and D. M. Treger, *Science* **294**, 1488 (2001).
- ³ D. D. Awschalom, N. Samarth, and D. Loss, *Semiconductor Spintronics and Quantum Computation* (Springer, Berlin, 2002).
- ⁴ T. C. Damen, L. Vina, J. E. Cunningham, J. Shah, and L. J. Sham, *Phys. Rev. Lett.* **67**, 3432 (1991).
- ⁵ J. Wagner, H. Schneider, D. Richards, A. Fischer, and K. Ploog, *Phys. Rev. B* **47**, 4786 (1993).
- ⁶ J. J. Baumberg, D. D. Awschalom, N. Samarth, H. Luo, and J. K. Furdyna, *Phys. Rev. Lett.* **72**, 717 (1994).
- ⁷ J. J. Baumberg, S. A. Crooker, D. D. Awschalom, N. Samarth, H. Luo, and J. K. Furdyna, *Phys. Rev. B* **50**, 7689 (1994).
- ⁸ A. P. Heberle, W. W. Rühle, and K. Ploog, *Phys. Rev. Lett.* **72**, 3887 (1994).
- ⁹ C. Buss, R. Frey, C. Flytzanis, and J. Cibert, *Solid State Commun.* **94**, 543 (1995).
- ¹⁰ S. A. Crooker, J. J. Baumberg, F. Flack, N. Samarth, and D. D. Awschalom, *Phys. Rev. Lett.* **77**, 2814 (1996).
- ¹¹ S. A. Crooker, D. D. Awschalom, J. J. Baumberg, F. Flack, and N. Samarth, *Phys. Rev. B* **56**, 7574 (1996).
- ¹² C. Buss, R. Pankoke, P. Leisching, J. Cibert, R. Frey, and C. Flytzanis, *Phys. Rev. Lett.* **78**, 4123 (1997).
- ¹³ J. M. Kikkawa, I. P. Smorchkova, N. Samarth, and D. D. Awschalom, *Science* **277**, 1284 (1997).
- ¹⁴ J. M. Kikkawa and D. D. Awschalom, *Nature* **397**, 139 (1998).
- ¹⁵ J. M. Kikkawa and D. D. Awschalom, *Phys. Rev. Lett.* **80**, 4313 (1998).
- ¹⁶ H. Ohno, *Science* **281**, 951 (1998).
- ¹⁷ Y. Ohno, R. Terauchi, T. Adachi, F. Matsukura, and H. Ohno, *Phys. Rev. Lett.* **83**, 4196 (1999).
- ¹⁸ R. I. Dzhioev, B. P. Zakharchenya, V. L. Korenev, D. Gammon, and D. S. Katzer, *JETP Lett.* **74**, 182 (2001).
- ¹⁹ R. I. Dzhioev, K. V. Kavokin, V. L. Korenev, M. V. Lazarev, B. Y. Meltser, M. N. Stepanova, B. P. Zakharchenya, D. Gammon, and D. S. Katzer, *Phys. Rev. B* **66**, 245204 (2002).
- ²⁰ Y. Yafet, *Phys. Rev.* **85**, 478 (1952).
- ²¹ R. J. Elliot, *Phys. Rev.* **96**, 266 (1954).
- ²² M. I. D'yakonov and V. I. Perel', *Zh. Eksp. Teor. Fiz.* **60**, 1954 (1971), [*Sov. Phys.-JETP* **33**, 1053 (1971)].

- ²³ G. L. Bir, A. G. Aronov, and G. E. Pikus, *Zh. Eksp. Teor. Fiz.* **69**, 1382 (1975), [*Sov. Phys.-JETP* **42**, 705 (1975)].
- ²⁴ F. Meier and B. P. Zakharchenya (Eds), *Optical Orientation* (North-Holland, Amsterdam, 1984).
- ²⁵ M. W. Wu and C. Z. Ning, *phys. stat. sol. (b)* **222**, 523 (2000).
- ²⁶ M. W. Wu and M. Kuwata-Gonokami, *Solid State Commun.* **121**, 509 (2002).
- ²⁷ M. W. Wu, *J. Phys. Soc. Jap.* **70**, 2195 (2001).
- ²⁸ M. W. Wu and C. Z. Ning, *Eur. Phys. J. B.* **18**, 373 (2000).
- ²⁹ M. W. Wu, *J. Supercond.: Incorporating Novel Mechanism* **14**, 245 (2001), cond-mat/0109258.
- ³⁰ F. X. Bronold, I. Martin, A. Saxena, and D. L. Smith, *Phys. Rev. B* **66**, 233206 (2002).
- ³¹ M. Q. Weng and M. W. Wu, *Phys. Rev. B* **66**, 235109 (2002).
- ³² M. Q. Weng and M. W. Wu, *J. Appl. Phys.* **93**, 410 (2003).
- ³³ M. Q. Weng and M. W. Wu, cond-mat/0302330; *Phys. Rev. B*, in press.
- ³⁴ M. Q. Weng and M. W. Wu, cond-mat/0210313; To be published in *phys. stat. sol. (b)*.
- ³⁵ M. Q. Weng and M. W. Wu, cond-mat/0303169.
- ³⁶ M. W. Wu and H. Metiu, *Phys. Rev. B* **61**, 2945 (2000).
- ³⁷ R. Eppenga and M. F. H. Schuurmans, *Phys. Rev. B* **37**, 10923 (1988).
- ³⁸ E. L. Ivchenko and G. E. Pikus, *Superlattices and Other Heterostructures* (Springer, Berlin, 1995).
- ³⁹ A. G. Aronov, G. E. Pikus, and A. N. Titkov, *Zh. Eksp. Teor. Fiz.* **84**, 1170 (1983), [*Sov. Phys.-JETP* **57**, 680 (1983)].
- ⁴⁰ H. Haug and A. P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors* (Springer-Verlag, Berlin, 1996).
- ⁴¹ G. D. Mahan, *Many-particle Physics* (Plenum, New York, 1981).
- ⁴² T. Kuhn and F. Rossi, *Phys. Rev. Lett.* **69**, 977 (1992).
- ⁴³ N. S. Averkiev, L. E. Golub, and M. Willander, *J. Phys.: Cond. Mat.* **14**, R271 (2002).
- ⁴⁴ W. H. Lau, J. T. Olesberg, and M. E. Flatté, *Phys. Rev. B* **64**, 161301 (2001).
- ⁴⁵ A. Malinowski, R. S. Britton, T. Grevatt, R. T. Harley, D. A. Ritchie, and M. Y. Simmons, *Phys. Rev. B* **62**, 13034 (2000).
- ⁴⁶ O. Madelung, M. Schultz, and H. Weiss (eds.), *Numerical Data and Functional Relationships in Science and Technology, Landolt-Börnstein, New Series*, vol. 17 (Springer-Verlag, Berlin, 1982).